


Flu Information: Guidance for Families Caring for a Child or Adult with Flu or Flu-Like Illness at Home



How Flu Spreads: The main way Flu viruses spread is from person to person in respiratory droplets of coughs and sneezes. The droplets from the infected person travel through the air and deposit on the mouth or nose of people nearby. The flu virus may also spread when a person touches respiratory droplets deposited on another person or object and then the person touches their own mouth, nose or eyes.



Stopping the Spread: To protect family members that are not sick:

- keep the sick person away from others, preferably in a room separate from common areas of the house; wear a face-mask - if available & tolerable -when sharing common spaces in the household
- remind sick person to cover coughs and clean their hands often using soap & water or alcohol-based hand rub
- all family members clean their hands often
- keep towels and wash cloths used by ill person separate from those used by other family members and clean bathroom surfaces with household disinfectant at least daily
- have only one person provide care to the sick person, if possible
- check with your health care provider to see if some persons in your home should take antiviral medications to prevent the flu
- have care giver avoid being face-to-face with the sick person



Monitor Household Members for Flu Symptoms:

Use the following check list to decide if you or someone in your household may have the flu:

- | | | |
|------------------------------|-----------------------------|--|
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Do you or your child have a cough? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Do you or your child have a sore throat? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Do you or your child have a headache, body aches, fatigue or chills? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Do you or your child have vomiting or diarrhea? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | Do you or your child have a fever of 100 degrees or higher? |

DO THE SYMPTOMS INDICATE FLU-LIKE ILLNESS?

If you checked **yes** to fever of 100 degrees or higher **AND** cough or sore throat you have an influenza-like illness. To avoid spreading the illness to others, persons with these symptoms should stay home, for at least 24 hours after their fever is gone, except to get medical care or for other necessities (without the use of fever-reducing medicine.) For persons working in health care and children younger than 5 years old the exclusion period should be continued for 7 days from symptom onset or until the resolution of symptoms, whichever is longer.

When to Consult a Doctor: warning signs that need medical attention

CHILDREN Warning Signs:

- Fast breathing or trouble breathing
- Bluish or gray skin
- Not drinking enough fluids
- Severe or persistent vomiting
- Not waking up or not interacting
- Being so irritable that child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough

ADULTS Warning Signs:

- Fast breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion
- Severe or persistent vomiting
- Flu-like symptoms improve but then return with fever and worse cough